

CSPE protocol

Adopted: 5/02 Revised: 7/16, 10/09

Abbreviations

The following is a list of abbreviations used at the University of Western States health centers. This list is not intended to be absolute nor universally comprehensive. Other standard abbreviations may be used that are not represented here. This list was drawn from a seed article [Hansen DT. Standard chart abbreviations in chiropractic practice. Chiropractic Technique 1991;3(2):96-103] and then edited. The CSPE committee deleted abbreviations that were uncommonly seen or used in our clinics as well as those felt to be so common that inclusion was unnecessary. Abbreviations were then added to the list by the committee as well as by other faculty on campus.

Abbreviations that have proven useful in our clinic system, but which may not be standard in a wider arena, have been placed in italics.

Note: Some symbols are encircled (e.g., pain = \bigcirc) which is appropriate for paper charting. In electronic charting the circle is usually omitted (e.g., pain = P).

This document is organized two ways: 1) organized by abbreviation and 2) organized by meaning.

Copyright © 2002 Western States Chiropractic College

2016 & 2009 Revision

Ronald LeFebvre, DC

Clincial Standards, Protocols & Education (CSPE) Committee

- Elizabeth Dunlop, DC
- Lorraine Ginter, DC
- Ronald LeFebvre, DC
- Ravid Raphael, DC

Other reviewers/contributors

- Joel Agresta, PT, DC (instructor for PT I and PT II)
- Lee McCaffrey, DC (instructor for Soft Tissue I and Rehab II)
- Karen E. Petzing, DC
- Anita Roberts, DC

Editorial assistant

Linda Myers

ABBREVIATIONS

ORGANIZED BY ABBREVIATION

Abbreviation	Definition
Ø	none
#	pound or fracture
P	pain
\uparrow	increased
 ↑↑	much increased
/d	per day
/m	per month
/wk	per week
/ WK	hours
"	minutes
<	less than
> ≈	greater than
	approximately
®	right
Θ	negative, normal, absent
Δ	change
μ	micro
\downarrow	decreased
$\downarrow\downarrow$	much decreased
1°	primary, first degree
1x/w	1 time per week
2°	secondary, second degree
2-4x/m	2-4 times per month
2xw/3w	2 times per week for 3 weeks
3°	tertiary, third degree
AAROM	active assisted range of motion
abd	abduction
abn	abnormal
add	adduction
adj	adjustment
ADL	activities of daily living
alt h/c	alternating hot/cold (contrast hot/cold)
am	morning
ant	anterior
AP	anterior-posterior
AROM	active range of motion
ASAP	as soon as possible
ASIS	anterior superior iliac spine
asym	asymmetrical
В	burning
B/B	bowel or bladder
B4	before
bi/bilat	bilaterally
bid	twice a day
BMR	basal metabolic rate

Abbreviation	Definition
BP or Bp	blood pressure
5	with
c/o	complains of
C/T	cervicothoracic
CA	cancer
CAD	coronary artery disease
CAD	cervical acceleration deceleration
СС	chief complaint
CFM	cross friction massage
CHF	congestive heart failure
chr	chronic
CMT	chiropractic manipulative therapy
COPD	chromic obstructive pulmonary disease
CR	contract relax
CRAC	contract relax antagonist contract
crep	crepitation(s)
cryo	cryotherapy
CT	Computed tomography
ctx	cervical traction
CVA	cerebrovascular accident
CVJ	
	costovertebral joint
Cx	
D	dull
d/t	due to
dbl	double
def	deficiency
delt	deltoid
dev	deviation
dim	diminished
DJD	degenerative joint disease
DOB	date of birth
DOD	date of draw
DOE	date of exam
DOI	date of injury
DOMS	delayed onset muscle soreness
DOT	department of transportation
DPT	diphtheria, pertussis, tetanus
DTR	deep tendon reflexes
Dx	diagnosis
e	electrical
EAM	external auditory meatus
ECG	electrocardiogram
EEG	electroencephalogram
EENT	ear, eyes, nose, and throat
EHL	extensor hallicus longus

	Definition
ENT	ear, nose and throat
ER	emergency room
eval	evaluation (-ed)
eve	evening
exac	exacerbation (-ed)
ext	external
Ext	extension
FH or FHx	family history
fix	fixation
flex or Flx	flexion
freq	frequency
FU	follow-up
Fx	fracture
GHJ	glenohumeral joint
GI	gastrointestinal
gmax	gluteus maximus
gmed	gluteus medius
gmin	gluteus minimus
grad	gradually
GTT	glucose tolerance test
GU	genitourinary
gyn	gynecological
h	hour
H&P	history and physical
НА	headache
Ham	hamstring
hern	herniation
HLA	human leukocyte antigen A (HLA B-27)
HMP	hot moist pack
HNP	herniated nucleus pulposis
hp	hot pack
HR	hold relax
HTN	high blood pressure/hypertension
HVG	high volt galvanism
HVT	high volt therapy
hx	history
Hz	hertz
IASTM	Instrument assisted soft tissue manipulation
IFC	interferential current
IIC	intermittent ischemic compression
Insp	inspiration
int	intermittent
int comp	Intermittent compression
ipsi	ipsilateral
IR or Infrd	infrared (lamp)
IU	International unit
IVD	intervertebral disc
jt	joint

Abbreviation	Definition
L	left
L/S	lumbosacral
L5	fifth lumbar
Lab	laboratory
Lat	lateral
LB	low back
LBP	low back pain
LE	lower extremity
LFAC	sine-low frequency alternating current
Lig	ligament
LLF	left lateral flexion
LMP	last menstural period
LMT	licensed massage therapist
LOC	loss of consciousness
LR, LROT	left rotation
LRR	left rotation restriction
LVG	low volt galvanic
Lx	lumbar
lx tr or lxt	lumbar traction
lxt or lx tr	lumbar traction
mA	milli amp
manip	manipulate
MED	minimal erythemal dose
meds	medications
MENS	microcurrent electrical nerve stimulation
MFTP	myofascial trigger point(s)
MI	myocardial infarction
min	minute
mm or msl	muscles
MMI	maximum (medical) improvement
MMR	measles, mumps, rubella
mo	month
mob	mobilize
mod	moderate
MP	motion palpation
MR	myofascial release
MRI	magnetic resonance imaging
MVA	motor vehicle accident
mVAS	mechanical visual analog scale
Ν	normal or negative
N & V	nausea & vomiting
n/a or NA	not applicable
N/C	no charge
NAD	no apparent distress
NARE	no apparent residual effects
NB	(nota bene) note well
NC	no change, non contributory
Neg	negative
NP	new patients, not performed

Abbreviation	Definition
NRE	no residual effects
NTT	normal (temperature) to touch
NTW	numbness tingling or weakness
0/0	on and off
OA	osteoarthritis
ob	obstetrics
occ	occasional; occiput
OPS	oral pain scale
OTC	over the counter drugs
OV	office visit
р	para
PA	posterior-anterior
Pafn	paraffin bath
palp	palpation
palp (P)	palpatory pain
Pass	passive
PE	physical examination
pec	pectoralis
PERRLA	pupils equal round regular react to light accommodate
perc	percussion
pfs	post-facilitation stretch
PHx	past history
PI	personal injury
PID	pelvic inflammatory disease
PIR	post isometric relaxation
piri	piriformis
pm	afternoon
Pmaj	pectoralis major
PMS	premenstrual syndrome
Pn	pain
PNF	proprioceptive neuromuscular facilitation
pos	positive
post	posterior
рр	patient presents
pps	pulses per second
PR	pulse & respiration or progress report
preg	pregnant
PRN	patient to return as needed
prod	produced
prog/px	prognosis
PROM	passive range of motion
prox	proximal
PS	pubic symphisis
PSIS	posterior superior iliac spine
PT	physiological therapeutics
pt	patient
PTR	patient to return
q	every
<u>Ч</u>	

QA r/o rad rads rec ref rehab resp rhom RLFR ROM ROT RR	quality assurancerule outradiatingradiationsrecommendrefer (-red)rehabilitationrespirationrhomboidright lateral flexion restrictionrange of motionrotation
rad rads rec ref rehab resp rhom RLFR ROM ROT	radiating radiations recommend refer (-red) rehabilitation respiration rhomboid right lateral flexion restriction range of motion rotation
rads rec ref rehab resp rhom RLFR ROM ROT	radiations recommend refer (-red) rehabilitation respiration rhomboid right lateral flexion restriction range of motion rotation
rec ref rehab resp rhom RLFR ROM ROT	recommend refer (-red) rehabilitation respiration rhomboid right lateral flexion restriction range of motion rotation
ref rehab resp rhom RLFR ROM ROT	refer (-red) rehabilitation respiration rhomboid right lateral flexion restriction range of motion rotation
rehab resp rhom RLFR ROM ROT	rehabilitation respiration rhomboid right lateral flexion restriction range of motion rotation
resp rhom RLFR ROM ROT	respiration rhomboid right lateral flexion restriction range of motion rotation
rhom RLFR ROM ROT	rhomboid right lateral flexion restriction range of motion rotation
RLFR ROM ROT	right lateral flexion restriction range of motion rotation
ROM ROT	range of motion rotation
ROT	rotation
RR	
	rotation restriction
RROT, RR	right rotation
RRR	right rotation restriction
RTW	return to work
Russ Stim	Russian stimulation
Rx	prescription
S	without (sans)
S	sharp
sac	sacrum
SCJ	sternoclavicular joint
SCM	sternocleido mastoid
seg	segment
sev	severe
SI	sacroiliac
sl	slight
SLR	straight leg raise
SMT	spinal manipulative therapy OR sensory motor
SOAAP	training Sub, Obj, Asses, Action, Plan
SOB	shortness of breath
SOL	space occupying lesion
sp	spinous process
spondy	spondylolisthesis
spr	sprain
SSLR	sitting straight leg raise
stab	stabilization
STE	self testicular examination
stim	stimulation
STM	soft tissue manipulation
str	strain
sup	supine
sw	swelling, swollen
SWD	short wave diathermy
Svid	-
	symptoms
sym T	symmetrical
•	tenderness taut and tender
T&T	

Abbreviation	Definition	
T&A	tonsils & adenoids (-ectomy)	
TENS	transcutaneous electrical nerve stimulation	
TFL	tensor fascia lata	
THER EX	therapeutic exercise	
TIA	transient ischemic attack	
TIA	transient ischemic attack	
tid	three times a day	
TOS	thoracic outlet syndrome	
TP	transverse process	
TPR	temperature, pulse, respiration	
TPT	trigger point therapy	
trap	trapezius	
troch	trochanter	
Tx	thoracic	
Tx or tx	treatment	
u/c	microcurrent	
UE	upper extremity	
uln	ulnar	
URI	upper respiratory infection	
US or u/s	ultrasound	
UTI	urinary tract infection	
UV	ultraviolet light therapy	

Abbreviation	Definition
Vag	vagina
val	valgus
var	varus
VAS	visual analog scale
VBI	vertebral basalar insufficiency
vert	vertebral
w/	with
w/cm2	watts per centimeter squared
w/o	without (sans)
WC, W/C, w/c	worker's compensation
wk	week
WNL	within normal limits
wp	whirlpool
wt	weight
х	times
у/о	year old
yest	yesterday

ORGANIZED BY MEANING

Definition	Abbreviation
1 time per week	1x/w
2 times per week for 3 weeks	2xw/3w
2-4 times per month	2-4x/m
abduction	abd
abnormal	abn
absent	Θ
active assisted range of motion	AAROM
active range of motion	AROM
activities of daily living	ADL
adduction	add
adjustment	adj
afternoon	pm
alternating hot/cold	alt h/c
anterior	ant
anterior superior iliac spine	ASIS
anterior-posterior	AP
approximately	~
as soon as possible	ASAP
asymmetrical	asym
basal metabolic rate	BMR
before	B4

Definition	Abbreviation
bilaterally	bi/bilat
blood pressure	BP or Bp
bowel or bladder	Ъ/В
burning	В
cancer	CA
cerebrovascular accident	CVA
cervical	Cx
cervical acceleration deceleration	CAD
cervical traction	ctx
cervicothoracic	C/T
change	Δ
chief complaint	СС
chiropractic manipulative therapy	CMT
chromic obstructive pulmonary disease	COPD
chronic	chr
complains of	c/o
computed tomography	СТ
congestive heart failure	CHF
contract relax	CR
contract relax antagonist contract	CRAC
3	

Definition	Abbreviation
coronary artery disease	CAD
costovertebral joint	CVJ
crepitation(s)	crep
cross friction massage	CFM
cryotherapy	cryo
date of birth	DOB
date of draw	DOD
date of exam	DOE
date of injury	DOI
decreased	\downarrow
deep tendon reflexes	DTR
deficiency	def
degenerative joint disease	DJD
delayed onset muscle soreness	DOMS
deltoid	delt
department of transportation	DOT
deviation	dev
diagnosis	Dx
diminished	dim
diphtheria, pertussis, tetanus	DPT
double	dbl
due to	d/t
dull	D
ear, eyes, nose, and throat	EENT
ear, nose and throat	ENT
electrical	е
electrocardiogram	ECG
electroencephalogram	EEG
emergency room	ER
evaluation (-ed)	eval
evening	eve
every	q
exacerbation (-ed)	exac
extension	Ext
extensor hallicus longus	EHL
external	ext
external auditory meatus	EAM
family history	FH or FHx
fifth lumbar	L5
first degree	1°
fixation	fix
flexion	flex or Flx
follow-up	FU
fracture	Fx or #
frequency	freq
gastrointestinal	GI
genitourinary	GU
glenohumeral joint	GHJ
glucose tolerance test	GTT
gluteus maximus	gmax

Definition	Abbreviation
gluteus medius	gmed
gluteus minimus	gmin
gradually	grad
greater than	>
gynecological	gyn
hamstring	Ham
headache	НА
herniated nucleus pulposis	HNP
herniation	hern
hertz	Hz
high blood pressure	HTN
high volt galvanism	HVG
high volt therapy	HVT
history	hx
history and physical	H&P
hold relax	HR
hot moist pack	HMP
hot pack	hp
hour	h
hours	6
human leukocyte antigen A (HLA B-27)	HLA
hypertension	HTN
increased	\uparrow
infrared (lamp)	IR or Infrd
inspiration	Insp
Instrument assisted soft tissue manipulation	IASTM
interferential current	IFC
intermittent	int
Intermittent compression	int comp
intermittent ischemic compression	lic
International unit	IU
intervertebral disc	IVD
ipsilateral	ipsi
joint	jt
laboratory	Lab
last menstural period	LMP
lateral	Lat
left	L
left lateral flexion	LLF
left rotation	LR, LROT
left rotation restriction	LRR
less than	<
licensed massage therapist	LMT
ligament	Lig
loss of consciousness	LOC
low back	LB
low back pain	LBP
low volt galvanic	LVG
lower extremity	LE

ABBREVIATIONS

lumbar Lx lumbar traction lxt or lxt tr lumbosacral Lx or lxt lumbosacral L/S manpetic resonance imaging MRI maximum (medical) improvement MMI mechanical visual analog scale mVAS medications meds micro μ microcurrent electrical nerve stimulation MENS minute min minutes " mobilize mod motor vehicle accident MVA motor vehicle accident MVA much increased 11 myocardial infarction MI myofascial release MRR myofascial release NA myofascial release NA motor vehicle accident MVA much increased 11 myofascial release MR myofascial release NR myofascial release NAD no apparent distress NAD no apparent distress NAD	Definition	Abbreviation
lumbar traction lxt or lxt r lumbor traction lx tr or lxt lumbosacral L/S magnetic resonance imaging MRI manipulate manip maximum (medical) improvement MMR mechanical visual analog scale mVAS medications meds micro μ microcurrent u/c microcurrent electrical nerve stimulation MENS minimal erythemal dose MED minute moh monting am moderate mod motion palpation MP much decreased ↓↓ much decreased ↓↓ much decreased MR myofascial release MR myofascial release MR myofascial release MR moderate Ne motor vehicle accident MI myofascial release MR myofascial release MR moderate Neg or Θ new pap		
lumbar traction lx tr or lxt lumbosacral L/S magnetic resonance imaging MRI manipulate manip maximum (medical) improvement MMI measles, mumps, rubella MMR mechanical visual analog scale mVAS medications meds micro μ microcurrent u/c microcurrent electrical nerve stimulation MENS milli amp mA minute min minutes " mobilize mod moderate mod motion palpation MP much decreased ↓↓ much increased ↑↑ muscles MR myofascial release MR myofascial release MR myofascial release NP no apparent distress NAD no apparent residual effects NARE no change, non contributory NC no residual effects NRE non		
Imboseral L/S magnetic resonance imaging MRI manipulate manip maximum (medical) improvement MMI measles, mumps, rubella MMR mechanical visual analog scale mVAS medications meds micro μ microcurrent electrical nerve stimulation MENS mili amp mA minimal erythemal dose MED minute mod moderate mod moting am motin palpation MP much decreased ↓↓ much increased MYA much increased MYA myofascial release MR myofascial release MR myofascial release NAP no apparent distress NAD no apparent distress NAD no apparent distress NAC no apparent distress NARE no charge N/C no residual effects NRE no c		
magnetic resonance imagingMRImanipulatemanipmaximum (medical) improvementMMImeasles, mumps, rubellaMMRmechanical visual analog scalemVASmedicationsmedismicroµmicrocurrentu/cmicrocurrent electrical nerve stimulationMENSmilli ampmAminuteminminutes"mobilizemobmotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓myofascial releaseMRmyofascial releaseMRmyofascial releaseMRmostersNAno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno change, non contributoryNCnorNTTnormalNor Gnormal (temperature) to touchNTTnormalNor Gnot applicablen/a or NAnot efformedNPnot contributoryNCno change, non contributoryNCno residual effectsNREnormal (temperature) to touchNTTnormalNor Gnot efformedNPnot efformedNPnot contributoryNCno contributoryNCno contributoryNCno contributoryNCnormal (temperature) to touchNTTnormalNor Gnot efformed		
manipulate manip maximum (medical) improvement MMI measles, mumps, rubella MMR mechanical visual analog scale mVAS medications meds micro µ microcurrent u/c microcurrent electrical nerve stimulation MENS miliamp mA minute min minutes " mobilize mob motor # motion palpation MP motor vehicle accident MVA much increased ↑↑ muscles mm or msl myofascial release MR myofascial release MR mostage vomiting N & V neagative Neg or Θ new patients NP no apparent distress NAD no apparent residual effects NARE no change, non contributory NC no residual effects NRE no charge N/C no residual effects<		
maximum (medical) improvement MMI measles, mumps, rubella MMR mechanical visual analog scale mVAS medications meds micro µ microcurrent u/c microcurrent electrical nerve stimulation MENS milli amp mA minimal erythemal dose MED minute min minute min mobilize mob moderate mod motion palpation MP much decreased ↓↓ much decreased ↓↓ much decreased ↓↓ much increased ↑↑ muscles mm or msl myofascial release MR myofascial release NAP no apparent distress NAD no apparent distress NAD no apparent residual effects NRE no change, non contributory NC nor esidual effects NRE nor contributory NC no		
measles, mumps, rubella MMR mechanical visual analog scale mVAS medications meds micro μ microcurrent u/c microcurrent electrical nerve stimulation MENS mili amp mA minimal erythemal dose MED minute min minute mob mobilize mob moderate mod motion palpation MP much decreased ↓↓ much decreased ↓↓ myofascial release MR myofascial release MR myofascial release MAD no apparent distress NAD no apparent residual effects NARE no change, non contributory NC nor NTT normal N or Θ not applicable n/a or NA not performed NP moderate MC moderate MC moderate MC <	-	
mechanical visual analog scalemWASmedicationsmedismicroμmicrocurrentu/cmicrocurrent electrical nerve stimulationMENSmilli ampmAminuteminminuteminmobilizemodmotoro palpationMPmotor vehicle accidentMVAmuch decreased↓↓myofascial releaseMRmyofascial releaseMRmodagaiveNPnausea & vomitingNFTPnausea & vomitingNAmyofascial releaseNAno apparent distressNADno apparent distressNAEno chargeN/Cno chargeNCno chargeNCno chargeNCno chargeNREnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (temperature) to touchNPnot applicablen/a or NAnot performedNPnot striging or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		
medications meds micro μ microcurrent u/c microcurrent electrical nerve stimulation MENS milli amp mA minute min minute min minutes " mobilize mod moderate mod motion palpation MP motor vehicle accident MVA much decreased ↓↓ myofascial release MR myofascial release MR myofascial release NA no apparent distress NARE no apparent distress NARE no change, non contributory NC nore Ø normal (temperature) to touch NTT normal Nor Θ not performed NP not performed NC nortices NT		
microμmicrocurrentu/cmicrocurrent electrical nerve stimulationMENSmili ampmAminual erythemal doseMEDminuteminminutes"mobilizemobmoderatemodmonthmomotion palpationMPmuch decreased↓↓much increased↑↑myofascial releaseMRRmyofascial releaseMRRmodapapationMFTPnausea & vomitingN & Vnotor vehicle accidentMImyofascial releaseMRRmyofascial releaseMRRmyofascial releaseNAREno apparent distressNADno apparent distressNADno chargeN/Cno residual effectsNREnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (temperature) to touchNTTnot applicablen/a or NAnot performedNPnot well (nota bene)NBnumbness tingling or weaknessNTWobstetricsobcoccasionaloccoffice visitOV		
microcurrentμ/cmicrocurrent electrical nerve stimulationMENSmili ampmAminimal erythemal doseMEDminuteminminutes"mobilizemobmoderatemodmonthmomotion palpationMPmuch decreased↓↓much increased↑↑myofascial releaseMRRmyofascial releaseMRRmotappationMFTPnausea & vomitingN & Vnoto relicit regressionNPnoto relicit regressionMRmuch increased↑↑musclesMRmyofascial releaseMRmyofascial releaseNAEno apparent distressNADno apparent distressNADno change, non contributoryNCnoreØnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (temperature) to touchNTnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsobcoccasionaloccoffice visitOV		meds
microcurrent electrical nerve stimulationMENSmili ampmAminimal erythemal doseMEDminuteminminutes"mobilizemobmoderatemodmonthmomotion palpationMPmuch decreased↓↓much increased↑↑myofascial releaseMRmyofascial releaseMRmyofascial releaseNAPno apparent distressNADno apparent distressNADno change, non contributoryNCno chargeN/Cnormal (temperature) to touchNTTnormalNor Θnormal (temperature) to touchNTTnormal (temperature) to touchNTUoccasionaloccoccisitOV		
milli ampmAminimal erythemal doseMEDminuteminminutes"mobilizemobmoderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreasedJJmusclesmm or mslmyofascial releaseMRmyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNADno apparent distressNADno chargeN/Cno charge, non contributoryNCnorØnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal not performedNPnot evell (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		
minimal erythemal doseMEDminuteminminutes"mobilizemobmoderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑myocardial infarctionMImyofascial releaseMRmyofascial releaseNAPno apparent distressNADno apparent distressNADno change, non contributoryNCnorMREnorØnormal (temperature) to touchNTTnormal (temperature) to touchNPnot applicablen/a or NAnot performedNPnot ewell (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciputoccocciputoccoffice visitOV		
minutesminminutes"mobilizemobmoderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑myocardial infarctionMImyofascial releaseMRmyofascial releaseNRmogativeNeg or Θnew patientsNADno apparent distressNADno chargeN/Cnon contributoryNCnorØnormal (temperature) to touchNTTnormal (temperature) to touchNTTnot performedNPnot evell (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciaputoccocciputoccocciputoccoffice visitOV		-
minutes"mobilizemobmoderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑myocardial infarctionMImyofascial releaseMRmyofascial releaseMRmyofascial releaseN& VnegativeNeg or Θnew patientsNADno apparent distressNAREno chargeN/Cno chargeN/Cnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (temperature) to touchNTTnot applicablen/a or NAnot performedNPnot settinging or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	minimal erythemal dose	
minutes"mobilizemobmoderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑musclesmm or mslmyofascial releaseMRmyofascial releaseMRmotagativeNeg or Θnew patientsNPno apparent distressNADno chargeN/Cno residual effectsNREnon contributoryNCnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (nota bene)NBnumbers tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		
moderatemodmonthmomorningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial rigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNPno apparent distressNAREno change, non contributoryNCno residual effectsNREnon contributoryNCnormal (temperature) to touchNTTnormal (temperature) to touchNTTnormal (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciputoccoffice visitOV		-
month mo morning am motion palpation MP motor vehicle accident MVA much decreased ↓↓ much increased ↑↑ muscles mm or msl myocardial infarction MI myofascial release MR myofascial trigger point(s) MFTP nausea & vomiting N & V negative Neg or Θ new patients NAD no apparent residual effects NARE no charge N/C no charge N/C non contributory NC nore Ø normal (temperature) to touch NTT normal (temperature) to touch NTT not applicable n/a or NA not performed NP note well (nota bene) NB numbness tingling or weakness NTW obstetrics ob occasional occ occiput occ	mobilize	mob
morningammotion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNPno apparent distressNADno change, non contributoryNCnoresidual effectsNREnon contributoryNCnoresidual effectsNREnormal (temperature) to touchNTTnormal (not applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccoffice visitOV	moderate	mod
motion palpationMPmotor vehicle accidentMVAmuch decreased↓↓much increased↑↑musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNPno apparent distressNADno change, non contributoryNCno chargeN/Cnon contributoryNCnoneØnormal (temperature) to touchNTTnormal (not applicablen/a or NAnot performedNPnot evell (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccoffice visitOV	month	mo
motor vehicle accidentMVAmuch decreased $\downarrow\downarrow$ much increased $\uparrow\uparrow$ musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θ new patientsNPno apparent distressNADno charge, non contributoryNCno chargeN/Cno contributoryNCno contributoryNCnoresidual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormal (not applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciputoccoffice visitOV	morning	am
much decreased $\downarrow\downarrow$ much increased $\uparrow\uparrow$ musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θ new patientsNPno apparent distressNADno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnoreØnormal (temperature) to touchNTTnormal not performedNPnot evell (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciputoccocciputoccoffice visitOV	motion palpation	MP
much increased↑↑musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNPno apparent distressNADno apparent residual effectsNAREno chargeN/Cno residual effectsNREnon contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnoreØnormal (temperature) to touchNTTnormal (temperature) to touchNPnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsobcoccasionaloccocciputoccoffice visitOV	motor vehicle accident	
musclesmm or mslmyocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Ønew patientsNPno apparent distressNADno dapparent residual effectsNAREno change, non contributoryNCno residual effectsNREnon contributoryNCno no contributoryNCno residual effectsNREnon contributoryNCnormal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsobcoccasionaloccocciputoccoffice visitOV	much decreased	
myocardial infarctionMImyofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Ønew patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnort applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoccocciputoccocciputoccoffice visitOV	much increased	$\uparrow \uparrow$
myofascial releaseMRmyofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Ønew patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormal not applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	muscles	mm or msl
myofascial trigger point(s)MFTPnausea & vomitingN & VnegativeNeg or Θnew patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnon contributoryNCno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormal not applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	myocardial infarction	MI
nausea & vomitingN & VnegativeNeg or Ønew patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	myofascial release	MR
negativeNeg or Θnew patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	myofascial trigger point(s)	MFTP
new patientsNPno apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsobcoccasionaloccocciputoccoffice visitOV	nausea & vomiting	N & V
no apparent distressNADno apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	negative	Neg or Θ
no apparent residual effectsNAREno change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocfice visitOV	new patients	NP
no change, non contributoryNCno chargeN/Cno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	no apparent distress	NAD
no chargeN/Cno residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	no apparent residual effects	NARE
no residual effectsNREnon contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	no change, non contributory	NC
non contributoryNCnoneØnormal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	no charge	N/C
noneØnormal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	no residual effects	NRE
normal (temperature) to touchNTTnormalN or Ønot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	non contributory	NC
normalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	·	Ø
normalN or Θnot applicablen/a or NAnot performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	normal (temperature) to touch	NTT
not performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		N or Θ
not performedNPnote well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV	not applicable	n/a or NA
note well (nota bene)NBnumbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		NP
numbness tingling or weaknessNTWobstetricsoboccasionaloccocciputoccoffice visitOV		NB
obstetricsoboccasionaloccocciputoccoffice visitOV		NTW
occasionaloccocciputoccoffice visitOV		ob
occiput occ office visit OV		
office visit OV		
	-	
	on and off	0/0

Definition	Abbreviation
oral pain scale	OPS
osteoarthritis	OA
over the counter drugs	OTC
pain	Pn
pain	®
palpation	Palp
palpatory pain	palp (P)
para	Р
paraffin bath	Pafn
passive	Pass
passive range of motion	PROM
past history	PHx
patient	Pt
patient presents	Рр
patient to return	PTR
patient to return as needed	PRN
pectoralis	Pec
pectoralis major	Pmaj
pelvic inflammatory disease	PID
per day	/d
per month	/m
per week	/wk
percussion	Perc
personal injury	PI
physical examination	PE
physiological therapeutics	PT
piriformis	Piri
positive	Pos
post isometric relaxation	PIR
posterior	Post
posterior superior iliac spine	PSIS
posterior-anterior	PA
post-facilitation stretch	Pfs
pound or fracture	#
pregnant	Preg
premenstrual syndrome	PMS
prescription	Rx
primary	1°
produced	Prod
prognosis	prog/px
progress report	PR
proprioceptive neuromuscular facilitation	PNF
proximal	Prox
pubic symphysis	PS
pulse, respiration or Progress Report	PR
pulses per second	Pps
pupils equal round regular react to light	PERRLA
accommodate	
quality assurance	QA
radiating	Rad

ABBREVIATIONS

Definition	Abbreviation
radiations	Rads
range of motion	ROM
recommend	Rec
refer (-red)	Ref
rehabilitation	Rehab
respiration	resp
return to work	RTW
rhomboid	rhom
right	®
right lateral flexion restriction	RLFR
right rotation	RROT, RR
right rotation restriction	RRR
rotation	ROT
rotation restriction	RR
rule out	r/o
Russian stimulation	Russ Stim
sacroiliac	SI
sacrum	sac
second degree	2°
secondary	2°
segment	seg
self testicular examination	STE
sensory motor training	SMT
severe	sev
sharp	S
short wave diathermy	SWD
shortness of breath	SOB
sine-low frequency alternating curr	LFAC
sitting straight leg raise	SSLR
slight	sl
soft tissue manipulation	STM
space occupying lesion	SOL
spinal manipulative therapy OR sensory motor training	SMT
spinous process	sp
spondylolisthesis	spondy
sprain	spr
stabilization	stab
sternoclavicular joint	SCJ
sternocleido mastoid	SCM
stimulation	stim
straight leg raise	SLR
strain	str
Sub, Obj, Asses., Action Plan	SOAP
supine	sup
swelling, swollen	SW
symmetrical	sym
symptoms	Sx
taut and tender	Т&Т
temperature, pulse, respiration	TPR

Definition	Abbreviation
tenderness	Т
tensor fascia lata	TFL
tertiary	3°
therapeutic exercise	THER EX
third degree	3°
thoracic	Tx
thoracic outlet syndrome	TOS
three times a day	tid
times	x
tonsils & adenoids (-ectomy)	T&A
transcutaneous electrical nerve stimulation	TENS
transient ischemic attack	TIA
transient ischemic attack	TIA
transverse process	TP
trapezius	trap
treatment	Tx or tx
trigger point therapy	TPT
trochanter	troch
twice a day	bid
ulnar	uln
ultrasound	US or u/s
ultraviolet light therapy	UV
upper extremity	UE
upper respiratory infection	URI
urinary tract infection	UTI
vagina	Vag
valgus	val
varus	var
vertebral	vert
vertebral basilar insufficiency	VBI
visual analog scale	VAS
watts per centimeter squared	w/cm2
week	wk
weight	wt
whirlpool	wp
with	с
with	w/
within normal limits	WNL
without (sans)	w/o
without (sans)	S
worker's compensation	WC, W/C, w/c
year old	у/о
yesterday	yest